

SENEGAL



General

Senegal - officially the Republic of Senegal - in West Africa is bordered by Mauritania in the North, Mali in the East, Guinea in the Southeast, and Guinea-Bissau in the Southwest. Senegal also borders Gambia, which separates Senegal's southern region of Casamance from the rest of the country. Senegal covers a land area of almost 19.7 Mha (million hectares) with in 2020 a population of 16.7 million, or 0.85 persons per ha (Wikipedia and United Nations, 2019).

Climate and geography

Source: esri

The climate in Senegal is Sahelian, but there is a rainy season. Mean tidal amplitudes are of the order of 1.0 to 1.5 m, with spring tide amplitudes up to 2.5 m. Sylla (1994) describes that in the Casamance River the tidal amplitude at the mouth is about 1.7 m to slightly over 0.5 m at Ziguinchor some 60 km upstream.

Marius (1982) describes that in Gambia and Senegal the areas with mangroves are concentrated along the banks and beaches in the estuaries of the Casamance, Gambia, Saloum, and Senegal rivers at a total area of about 500,000 ha (Figure 1). Parts of the mangrove areas have been replaced by *tannes*. *Tannes* are areas where the mangroves were cleared for paddy fields or salt pans. Sylla (1994) describes that rice growing in the mangrove zone started in the middle of the 18th century in Guinea and Sierra Leone. Traditional systems are still the most widespread and they are applied, for example in Senegal (diola system), Guinea, Guinea-Bissau (bolanha system) and Sierra Leone. The diola and bolanha systems consist of small basins or strips of land that are surrounded by dikes. Within these polders the rice is cultivated on ridges. Tidal drainage is required to flush the salts and acids that have accumulated in the polders during the dry season.




Figure 1. Mangrove area in Senegal and Gambia (Marius, 1982)

Since the early 1960s the soils in the mangrove areas have been studied, while after reclamation these soils became acid. Since 1972 systematic soil surveys have been conducted in the three estuaries, resulting in soil maps and relevant physiographic background information. Detailed maps have been made for specific reclamation projects.

The traditional systems of rice cultivation have functioned well until the persisting droughts started in 1969. The most affected zones are mainly in the northern, and drier coastal part of West Africa, including Senegal, Gambia, Guinea-Bissau and to a some extent Guinea.

Marius (1982) describes that the intertidal clay flats facilitated the construction of small polders. Dikes were 1.0 to 1.5 m high with a crest level of 0.20 m above maximum spring tide level. For sluices hollow tree trunks, preferably of palms, were embedded at various levels in the dike body, enabling the maintenance of various water levels in the polder. The lowest sluice was situated below the original surface with its outlet in the trench. For gates fibrous stops impregnated with clay were used and sometimes wooden flap gates. After clearing a grid of superficial ditches was dug and the excavated flat topsoil clods were laid on the interjacent surface areas to form raised cambered beds of a few square metres each. Impoldering normally started at the inland side of the Rhizophora zone and later expanded towards the river bank. In areas with erosive tidal currents the rice polders were protected by a series of peripheral polders. Moreover, outer dikes were constructed several tens of metres from the banks. Leaving a fringe of mangrove forest in the front land (Figure 2).




Figure 2. Traditional rice polders in the Casamance estuary (Marius, 1982)

Marius (1982) also mentions that even before 1972 large-scale reclamation projects for paddy production have been realised. The more important ones are situated in the Lower Casamance where the mangrove flats penetrate into the valleys of the tributaries. Upstream of the tidal areas the valleys were dammed to regulate the fresh water supply. Concrete dams with wooden gates in the riverbed exclude saline tides and enable increased drainage capacity. In some polders, reclaimed in 1967, after desalinisation and leaching acidification occurred.

In a report of the International Bank for Reconstruction and Development - International Development Association (1971) reference is made of seven pilot rice development schemes in the Casamance Region that started between 1967 and 1970. In each of them, the approach has been to introduce higher yielding varieties of paddy, to improve cultivation methods and to promote the use of draft oxen by means of a field extension service (Figure 3). In addition, in the case of swamp rice, drainage improvement works were carried out. Together, these projects covered an area of about 3,000 ha of paddy. The project would include the construction of 2000 ha of irrigation and drainage works in swampy areas.




Figure 3. Casamance Rice Project, proposed development area (International Bank for Reconstruction and Development - International Development Association, 1971)

In a World Bank report of 2013 reference is made of actions that would also include construction of intermediate storage and regulatory structures upstream, putting in place small dikes and drainage systems which would not require heavy investments, provision of internal access and crossing structures, and implementation of erosion control systems including agroforestry activities. The project would finance the necessary investments for the development and or rehabilitation of: i) flooded and floodable valleys for rice and aquaculture production; ii) horticulture perimeters. The selection of valleys such as Sofa Niama, Temento (Kolda), Séléky, Tengory and Suelle (Ziguinchor), Médina Souané (Sédhiou) would be based on specific criteria including areas cleared of landmines, absence of land disputes, existence of youth or women associations, technical feasibility, basic diking facilities (if necessary the dikes would be strengthened by refilling of materials), valleys equipped with salt control dams, and expected impacts. In rice valleys the project would target on lands that had basic dike facilities, but without gate-controlled spillways. This would also be implemented in valleys equipped with salt control dams and for which there were available data. The targeted regions were Ziguinchor and Sédhiou, and to a lesser extent the region of Kolda, based on the existing potential. Eligible activities included works related to rehabilitation of flooded and floodable valleys, and shallows.

Altogether the reports showed that there would be a larger polder area along the rivers, but it was difficult to quantify how large it really was.

Existing polders

In a World Bank report of 1972 the Dagana, Debi and Lampsar polders were mentioned (Figure 4). The project included the following: i) construction of an irrigation system in Dagana Polder permitting year-round cultivation of 2,730 ha; ii) improving water control in the Debi Polder and increasing the irrigated area from 260 ha to 1,025 ha; iii) improving water control in the seven ponds forming the Lampsar subproject and increasing the irrigated area from 880 ha to 1,080 ha; iv) strengthening the *Societe d'Amenagement et d'Exploitation des Terres du Delta (SAED)* with staff, facilities, and equipment to carry out the project and to provide support for farmers who use land developed under the project; v) constructing, equipping, and staffing a training centre at Dagana for SAED personnel and for farmers; vi) carrying out, through collaboration with *Institut de Recherches agronomiques tropicales (IRAT)*, a program of applied research to support agricultural operations at Dagana; vii) expanding the SAED farm machinery pool to permit of land preparation services to farmers using land developed under the project (World Bank, 1972). In the World Bank evaluation report (1979) mention is made of differences in project implementation, compared to the objectives. However, no significant differences in reclaimed areas could be identified.




Figure 4. The Senegal River Polders Project area (World Bank, 1972)

The Group Polder Development (1982) provided information on three polders. These are:

- in the framework of the Lower Casamance Project the experimental polders Medina (10 ha) and Dieba (30 ha) (Figure 5). The polders are located in the Delta of the Casamance River;
- the Richard Toll (Caser) Polder along the Senegal River. This is a polder of some thousand ha for the growth of sugar cane.

Moens and Wanders presented the following data on the area of several polders: Lampsar polder 10,000 ha, Dagana Polder 3,000 ha and Nianga Polder 1,000 ha.

In a report of the African Development Fund (2005) it is stated that the project will be implemented in lands in the Lower Casamance River area that have basic diking facilities, but without gate-controlled spillways. The dikes would be strengthened (refilling of materials, fixing with vetiver, etc.) and equipped with water control structures. This component would concern schemes undertaken in recent years under programme of the World Food Program (WFP) (several hundreds of kilometres of dikes), as well as future ones, and would protect about 4,500 ha of land.

Some characteristic data of the existing and proposed polders are shown in Table I.




Figure 5. Lower Casamance Project (Group Polder Development, 1982)

Proposed polders

At request of the Senegal River Basin Development Authority, under the title *Fight against weed pests in the Senegal River Delta*, the *Rijksdienst voor Ondernemend Nederland* (2010) invited proposals for the construction of eight polders along the Senegal River. These polders would include 63 km of polder dikes, about 60 km of drains, about 8 pumping systems and at least 8 inlet structures. Four of these polders would be constructed on the Senegal side of the river and four on the Mauritania side. The polders would be located close to the towns of Rosso and Richard Toll along existing agriculture fields. Another important aspect of the project would be the clearance of the Typha – a type of reet - and the preparation of the land in the new polders. The project proposed the creation of polders where the Typha cannot grow: a sustainable solution as it would mean that existing Typha would only have to be cleared once at the start of the project. As soon as the waterline would not be obstructed by Typha anymore, local inhabitants would have easier access to water. The polders would have to create new agricultural ground in which rice and other agricultural products such as vegetables can be grown. In the framework of this programme Royal HaskoningDHV (2013), has produced a master plan to develop the Senegal River Delta and designs for the eight polders with a total area between 3000 and 4000 ha as a pilot project. An artist impression is shown in Figure 6.



Figure 6. Artist impression of the proposed polders in the Senegal River Delta
(source: Royal HaskoningDHV, 2013)

References

- African Development Fund, 2005. Republic of Senegal. *The Casamance Rural Development Support Project (PADERCA)*. Appraisal Report.
- Beye, G., 1973. *Acidification of mangrove soils after empoldering in lower Casamance*. Effects of the type of reclamation system used. In: Dost, H. (ed.). Acid sulphate soils – Proceedings International Symposium. ILRI publ. 18, Vol. II, Wageningen, the Netherlands.
- Beye, G., 1973. *Bilan de cinq années d'études du dessalement des sols du polder de Medina (Basse Casamance)*. Inst. Rech. Agr. Trop. Djibelor (in French).
- Beye, G., 1973. Une méthode simple de dessalement des sols de tannes de Casamance: le paillage. *Agr. Trop.* 26: 537-549 (in French).
- Cadillac, H. 1965. *Étude agropédologique de quelques dépressions rizicultivables de la région de Bignona (Casemance)*. Orstrom, Dakar, Senegal. 80 p. (in French).
- Group Polder Development, Department of Civil Engineering, Delft University of Technology. 1982. *Polders of the World. Compendium of polder projects*. Delft, the Netherlands
- International Bank for Reconstruction and Development - International Development Association, 1971. *Casamance Rice Project, Senegal*.
- Marius, C., 1980. Les sols des mangroves et tannes de la region des Kalounayes. Orstrom, Dakar, Senegal (in French).
- Marius, C., 1982. Acid sulphate soils of the Mangrove area of Senegal and Gambia. In: Dost, H. and N. van Breemen (eds.). Proceedings of the Bangkok Symposium on Acid Sulphate soils. Bangkok, Thailand. International Institute for Land Reclamation and Improvement (ILRI), Wageningen, the Netherlands.
- Moens, A. and A.A. Wanders, 1983. *Landbouwmechanisatie en Landbouwwerktuigenindustrie in Mali*. Wageningen, the Netherlands (in Dutch).
- Rijksdienst voor Ondernemend Nederland, 2010. *Fight against weed pests in the Senegal River Delta*. <https://www.rvo.nl/subsidies-regelingen/projecten/fight-against-weed-pests-senegal-river-delta>
- Royal HaskoningDHV, 2013. *Tackle typha in Senegal River Delta. Polders along Senegal River*. Amersfoort, the Netherlands.
- Sylla, M., 1994. *Soil salinity and acidity: spatial variability and effects on rice production in West Africa's mangrove zone*. PhD thesis Wageningen University. Wageningen, the Netherlands.
- United Nations, Department of Economic and Social Affairs, Population Division. 2019. *World population prospects, medium prognosis. The 2019 revision*. New York, USA.
- Vieillefon, J., 1977. Les sols des mangroves et tannes de Basse Casamance. Mém. ORSTROM 83, 291 p. (in French).
- World Bank, 1972. Report and recommendation of the President to the Executive Directors on a proposed development credit to the Republic of Senegal for a river polders project. Report No. P-1157-SE.
- World Bank, 1979. PPAR, Senegal - River Polders Project (Credit 350-SE), OED Report No. 2777 dated December 26, 1979.
- World Bank, 1987. *Project performance audit report. Lake Chad Polders Project*. Operations Evaluation Department, Report no. 6751 (restricted availability). Washington DC, USA.
- World Bank, 2013. *Casamance Development Pole Project*. Project appraisal document.

Bart Schultz

Lelystad, January 2021

Table I. General characteristics of existing and proposed polders in Senegal

Name	Reclamation	Area in ha	Type *)	Latitudes	Longitudes	Elevation in m+MSL	Land use
<i>Existing polders</i>							
Medina Polder		10	RLL	12° 49' N	15° 58' W	23	
Dieba Polder		30	RLL	12° 50' N	16° 02' W	25	
Richard Toll Polder		1,000	RLL	16° 27' N	15° 43' W	6	
Dagana Polder		3,000	RLL	16° 31' N	15° 31' W	7	
Debi Polder		1,025	RLL	16° 28' N	16° 15' W	4	
Lampsar polders		10,000	RLL	16° 07' N	16° 21' W	4	
Nianga Polder		1,000	RLL	12° 42' N	15° 15' W	47	
Polders in the Lower Casamance River area		4,500	RLL	12° 26' N	16° 33' W	2	
Sub-total		20,565					
<i>Proposed polders</i>							
4 polders in Senegal River Delta							
Sub-total							
Total		20,565					

*) RLL = reclaimed low-lying land; LGS = land gained on the sea; DL = drained lake